

ANALLERGENIC



- Formulated with feather protein hydrolysate, a protein source containing exclusively free amino acids and very low molecular weight oligopeptides (proteins).



- Purified maize starch instead of whole cereals and a specially designed palatability enhancer; manufactured in a production line designed to minimise contamination.



- A combination of nicotinamide, inositol, choline, histidine and pantothenic acid helps reduce water losses through the skin and strengthen the barrier effect of the skin.



- The synergistic antioxidant complex helps neutralise free radicals.

COMPOSITION

Maize starch, feather hydrolysate with low molecular weight (source of L-amino acids and oligopeptides), copra oil, soya oil, minerals, vegetable fibres, chicory pulp, Fructo-Oligo-Saccharides (FOS), fish oil, mono- and diglycerides of fatty acids esterified with citric acid, animal fat, marigold extract (source of lutein).

Additives (kg) **

Nutritional additives:

Vitamin A: 30600IU, Vitamin D3: 800IU, E1 (Iron): 42mg, E2 (Iodine): 3.7mg, E4 (Copper): 13mg, E5 (Manganese): 55mg, E6 (Zinc): 198mg, E8 (Selenium): 0.1mg.
Preservatives - Antioxidants.



Available in
3kg and 8kg bag



Approximate size

KEY VALUES

per 100g
as fed

Protein	18g
Fat content	16.5g
Carbohydrate	44.2g
NFE	48g
Dietary fibre	6g
Crude fibre	2.2g
Omega 6	3.91g
Omega 3	0.6g
EPA+DHA	0.2g
Linoleic acid	3.65g
Calcium	1.01g
Phosphorus	0.8g
Sodium	1g
Metabolisable energy*	388kcal

*Calculated according to NRC2006

**NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.

CLINICAL TRIALS OR STUDIES CARRIED OUT WITH ROYAL CANIN ANALLERGENIC

- Mougeot I. & al. Clinical efficacy of a highly hydrolyzed poultry feather protein-based diet for canine AFR diagnosis and dietary management: a 12 case pilot study. Poster presented at WINSS 2013; October 1-4, Portland, Oregon (USA).
- Freiche V. & al. A Very Low Molecular Weight Poultry Feather Hydrolyzed-Based Extruded Diet Allows Idiopathic Canine IBD Clinical Management Without Immunosuppressive Treatment: A 13-Case Prospective Pilot Study. Short communication presented at ECVIM congress 2014; Sept 4-6, Mainz (Germany).

ADVERSE REACTION TO FOOD



SKIN

RECOMMENDED FOR DOGS WITH

- Adverse Food Reactions (AFR):
 - Food allergy
 - Food intolerance
 - With dermatological and/or gastro-intestinal signs
 - Food elimination trial
- Inflammatory Bowel Disease (IBD)

NOT RECOMMENDED FOR

- Pregnancy, lactation, growth
- Pancreatitis

RECOMMENDED DAILY INTAKE & FEEDING DURATION

Dog's weight	thin		normal		overweight	
	grams	cups	grams	cups	grams	cups
2kg	54	5/8	48	4/8	41	4/8
4kg	91	1	80	7/8	69	6/8
6kg	123	1 3/8	109	1 2/8	94	1
8kg	153	1 5/8	135	1 4/8	116	1 2/8
10kg	181	2	159	1 6/8	138	1 4/8
15kg	245	2 5/8	216	2 3/8	186	2
20kg	304	3 6/8	268	3	231	2 4/8
25kg	360	4	317	3 4/8	273	3
30kg	413	4 4/8	363	4	314	3 4/8
35kg	463	5 1/8	408	4 4/8	352	3 7/8
40kg	512	5 5/8	450	4 7/8	389	4 2/8
50kg	605	6 5/8	533	5 7/8	460	5
60kg	694	7 5/8	611	6 3/8	527	5 6/8
70kg	779	8 4/8	685	7 4/8	592	6 4/8
80kg	861	9 4/8	758	8 2/8	654	7 1/8

When a food allergy or a food intolerance is suspected, the dog's diet should be changed without any food transition. Nutritional management with Anallergenic may be continued for the pet's lifetime.

It is recommended to divide the daily intake into 2 meals