

# HYPOALLERGENIC



Available in  
400g can



- Soy protein hydrolysate, which is composed of low molecular weight peptides, is highly digestible and of very low allergenicity.



- A combination of nicotinamide, inositol, choline, histidine and pantothenic acid helps reduce trans-epidermal water loss and strengthens the barrier effect of the skin.

- Enriched with linoleic acid, EPA and DHA, essential and non-essential fatty acids known to modulate cutaneous reactions and help maintain skin health.
- The synergistic antioxidant complex helps neutralise free radicals.

## KEY VALUES

	per 100g as fed	100g dry matter
Protein	6g	23.5g
Fat content	3.5g	13.7g
Carbohydrate	11.1g	43.5g
NFE	12.4g	48.6g
Dietary fibre	3.3g	12.9g
Crude fibre	2g	7.8g
Omega 6	1.7g	6.7g
Omega 3	0.28g	1.1g
EPA+DHA	0.2g	0.8g
Calcium	0.19g	0.7g
Phosphorus	0.18g	0.7g
Sodium	0.2g	0.8g
Potassium	0.22g	0.9g
Magnesium	0.02g	0.1g
Metabolisable energy*	95kcal	372.5kcal

## COMPOSITION

Pea starch, soya protein hydrolysates, sunflower oil, cellulose, fish oil, minerals, hydrolysed chicken liver, lecithin, taurine, marigold meal (rich in lutein), trace elements (including chelated trace elements), vitamins.

## Additives (kg) \* \*

### Nutritional additives:

Vitamin A: 5000IU, Vitamin D3: 340IU, E1 (Iron): 25mg, E2 (Iodine): 2mg, E4 (Copper): 6mg, E5 (Manganese): 32 mg, E6 (Zinc): 120mg.

### Technological additives:

Pentasodium triphosphate: 3g.

\*Calculated according to NRC2006

**\*\*NOTE:** Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.

### CLINICAL TRIALS OR STUDIES CARRIED OUT WITH ROYAL CANIN HYPOALLERGENIC

- Biourge V. Diagnosis of adverse reactions to food in dogs: efficacy of a soy-isolate hydrolyzate-based diet. J Nutr 2004; 134: 2062S-2064S.
- Horvath-Ungerboeck C. & al. Efficacy of a hydrolysed diet in 80 pruritic dogs: a prospective study. Proceedings of the 27th annual ESVD-ECVD congress, Salzburg (Austria), 11-13 September 2014, p18.

# ADVERSE REACTION TO FOOD



SKIN

## RECOMMENDED FOR DOGS WITH

- Adverse Food Reactions (AFR):
  - Food allergy
  - Food intolerance
  - With dermatological and/or gastro-intestinal signs
  - Food elimination trial
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Chronic diarrhoea
- Small intestinal bacterial overgrowth (SIBO)

## NOT RECOMMENDED FOR

- Pregnancy, lactation, growth
- Pancreatitis

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

Dog's weight	thin		normal		overweight	
	grams	cans 400g	grams	cans 400g	grams	cans 400g
2kg	220	1/2	195	1/2	170	1/2
3kg	300	3/4	265	3/4	230	1/2
4kg	370	1	330	3/4	285	3/4
5kg	440	1	385	1	335	3/4
6kg	505	1 1/4	445	1	385	1
7kg	565	1 1/2	500	1 1/4	430	1
8kg	625	1 1/2	550	1 1/2	475	1 1/4
9kg	685	1 3/4	600	1 1/2	520	1 1/4
10kg	740	1 3/4	650	1 3/4	560	1 1/2
15kg	1005	2 1/2	885	2 1/4	760	2
20kg	1245	3	1095	2 3/4	945	2 1/4
25kg	1470	3 3/4	1295	3 1/4	1120	2 3/4
30kg	1685	4 1/4	1485	3 3/4	1280	3 1/4
35kg	1895	4 3/4	1665	4 1/4	1440	3 1/2
40kg	2095	5 1/4	1840	4 1/2	1590	4

When a food allergy or a food intolerance is suspected, the dog's diet should be changed without any food transition. The hypoallergenic nutritional management will be continued for the pet's lifetime.

1 can is equivalent to 95g of Hypoallergenic dry product.

A mixed feeding guideline can be found at the end of this section.

It is recommended to divide the daily intake into 2 meals.