SENSITIVITY CONTROL



 Duck and tapioca are the only sources of protein and carbohydrate in this diet.
Both are rarely involved in food allergy, reducing the risk of GI or dermatological reacions.



 A combination of nicotinamide,inositol, choline, histidine and pantothenic acid helps reduce trans-epidermal water loss and strengthens the barrier effect of the skin.



 Eicosapentaenoic and docosahexaenoic acids (omega-3 long chain fatty acids) modulate skin reactions and contribute to the intestinal mucosal integrity.



 Contains a combination of highly digestible proteins, prebiotics, beet pulp and fish oil to ensure maximum digestive security. This food is gluten and lactose-free.



Tapioca, dehydrated duck meat, hydrolysed poultry proteins, vegetable fibres, animal fats, beet pulp, fish oil, soya oil, psyllium husks and seeds, minerals, Fructo-Oligo-Saccharides (FOS), marigold extract (source of lutein).



Additives (kg) * *

Nutritional additives:

Vitamin A: 25000IU, Vitamin D3: 800IU, E1 (Iron): 41mg, E2 (Iodine): 2.8mg, E4 (Copper): 10mg, E5 (Manganese): 54mg, E6 (Zinc): 203mg, E8 (Selenium): 0.1mg.

Preservatives - Antioxidants.

**NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.







Approximate size


KEY VALUES	per 100g as fed		
Protein	21g		
Fat content	9g		
Carbohydrate			
NFE	48.9g		
Dietary fibre			
Crude fibre	4.4g		
Omega 6	1.98g		
Omega 3			
EPA+DHA			
Biotin	1.89g		
Linoleic acid			
Phosphorus			
Zinc	23.8mg		
Biotin			
Pantothenic acid	14.7mg		
Niacin	49mg		
Metabolisable energy*	334kcal		

*Calculated according to NRC2006

LITERATURE ON WHICH ROYAL CANIN SENSITIVITY CONTROL IS BASED

• Roudebush P. Ingredients and foods associated with adverse reactions in dogs and cats. Veterinary Dermatology 2013; 24(2): 293–294.





ADVERSE REACTION TO FOOD



RECOMMENDED FOR DOGS WITH

NOT RECOMMENDED FOR

- Adverse Food Reactions (AFR):
 - Food allergy
 - Food intolerance

• Acute pancreatitis (may be fed in cases with chronic pancreatitis)

RECOMMENDED DAILY INTAKE & FEEDING DURATION

Dog's weight	thin		normal		overweight	
	grams	cups	grams	cups	grams	cups
2kg	63	6/8	55	5/8	48	4/8
4kg	106	1 2/8	93	1	80	7/8
6kg	143	1 5/8	126	1 3/8	109	1 2/8
8kg	178	2	156	1 6/8	135	1 4/8
10kg	210	2 3/8	185	2 1/8	160	1 6/8
15kg	285	3 2/8	251	2 7/8	217	2 4/8
20kg	354	4	311	3 4/8	269	3
25kg	418	4 6/8	368	4 1/8	318	3 5/8
30kg	479	5 3 / 8	422	4 6/8	364	4 1/8
35kg	538	6	473	5 3/8	409	4 5/8
40kg	595	6 6/8	523	5 7/8	452	5 1/8
50kg	703	7 7/8	619	7	534	6
60kg	806	9 1/8	709	8	612	6 7/8
70kg	905	10 2/8	796	9	688	7 6/8
80kg	1000	11 2/8	880	9 7/8	760	8 4/8

When a food allergy or a food intolerance is suspected, the dog's diet should be changed without any food transition. The nutritional management may be continued for the pet's lifetime.

It is recommended to divide the daily intake into 2 meals.

FURTHER INFORMATION

Skin care is recommended for dogs with skin conditions not relating to diet.



