

SKIN CARE



- A combination of essential fatty acids from linseed, soya, borage and fish oils enhances the skin's natural strength and help reduce the signs of skin disease.



- A combination of nicotinamide, inositol, choline, histidine and pantothenic acid helps reduce trans-epidermal water loss and strengthens the barrier effect of the skin.



- Contains a combination of highly digestible proteins (L.I.P.* proteins), prebiotics, beet pulp, rice and fish oil to ensure maximum digestive security.



- The synergistic antioxidant complex helps neutralise free radicals.

COMPOSITION

Maize, wheat gluten*, rice, tapioca, animal fats, maize gluten, minerals, soya oil, beet pulp, fish oil, vegetable fibres, flax seed, hydrolysed animal proteins, Fructo-Oligo-Saccharides (FOS), mono- and diglycerides of fatty acids esterified with citric acid, borage oil, marigold extract (source of lutein).

Additives (kg)***

Nutritional additives:

Vitamin A: 30100IU, Vitamin D3: 800IU, E1 (Iron): 56mg, E2 (Iodine): 5.6mg, E4 (Copper): 11mg, E5 (Manganese): 73mg, E6 (Zinc): 220mg, E8 (Selenium): 0.13mg.

Preservatives - Antioxidants.



Available in 2kg, 8kg and 12kg bag



Approximate size

KEY VALUES

per 100g as fed

Protein	23g
Fat content	16g
Carbohydrate	38.4g
NFE	43.9g
Dietary fibre	7.4g
Crude fibre	1.9g
Omega 6	3.64g
Omega 3	1.04g
EPA+DHA	0.41g
Calcium	0.76g
Phosphorus	0.58g
Sodium	0.3g
Metabolisable energy**	390kcal

*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation

**Calculated according to NRC2006

***NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.

CLINICAL TRIALS OR STUDIES ON WHICH ROYAL CANIN SKIN CARE IS BASED

- Watson AL & al. Dietary constituents are able to play a beneficial role in canine epidermal barrier function. *Experimental Dermatology* 2006; 15: 74-81.
- Van Beeck FL & al. The effect of long term feeding of skin barrier-fortified diets on the owner-assessed incidence of atopic dermatitis symptoms in labrador retrievers. *J Nutr Sci.* 2015 Feb 12;4:e5.

DERMATOSIS AND HAIR LOSS



SKIN

RECOMMENDED FOR DOGS WITH

- Atopic dermatitis
- Flea Allergy Dermatitis (FAD)
- Pyodermatitis
- Otitis externa
- Skin healing
- Dermal function maintenance (dermatosis, hair loss)

NOT RECOMMENDED FOR

- Pregnancy, lactation, growth

RECOMMENDED DAILY INTAKE & FEEDING DURATION

Dog's weight	thin		normal		overweight	
	grams	cups	grams	cups	grams	cups
11kg	194	1 7/8	170	1 5/8	147	1 3/8
15kg	244	2 2/8	215	2	186	1 6/8
20kg	303	2 7/8	267	2 4/8	230	2 1/8
25kg	358	3 3/8	315	3	272	2 5/8
30kg	411	3 7/8	362	3 3/8	312	3
35kg	461	4 3/8	406	3 7/8	351	3 3/8
40kg	510	4 7/8	449	4 2/8	387	3 5/8
45kg	557	5 2/8	490	4 5/8	423	4
50kg	603	5 6/8	530	5	458	4 3/8
55kg	647	6 1/8	570	5 3/8	492	4 5/8
60kg	691	6 4/8	608	5 6/8	525	5
65kg	734	7	646	6 1/8	558	5 2/8
70kg	776	7 3/8	683	6 4/8	589	5 5/8
75kg	817	7 6/8	719	6 6/8	621	5 7/8
80kg	857	8 1/8	754	7 1/8	652	6 1/8
85kg	897	8 4/8	790	7 4/8	682	6 4/8
90kg	937	8 7/8	824	7 6/8	712	6 6/8
95kg	975	9 2/8	858	8 1/8	741	7
100kg	1014	9 5/8	892	8 4/8	770	7 2/8

As soon as the first signs of skin disease appear, the diet should be used for a 2-month period to obtain visible results. For chronic disease, Skin Care can be maintained for the dog's entire lifetime.

It is recommended to divide the daily intake into 2 meals.