

HYPOALLERGENIC



- Soy protein isolate digest is composed of low molecular weight peptides, so is highly digestible and of very low allergenicity.



- Very high amounts of biotin, niacin and pantothenic acid, in combination with the zinc-linoleic acid complex, reduce water losses through the skin and strengthen the barrier effect of the skin.



- Eicosapentaenoic and docosahexaenoic acids, omega-3 long chain fatty acids, modulate skin reactions and contribute to the intestinal mucosal integrity.



- Contains a combination of highly digestible proteins (L.I.P.* proteins), prebiotics, beet pulp, rice and fish oil to ensure maximum digestive security.

COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, vegetable fibres, minerals, hydrolysed poultry liver, soya oil, beet pulp, fish oil, Fructo-Oligo-Saccharides (FOS), borage oil, marigold extract (source of lutein).

Additives (kg)***

Nutritional additives:

Vitamin A: 25300IU, Vitamin D3: 800IU, E1 (Iron): 41mg, E2 (Iodine): 3mg, E4 (Copper): 10mg, E5 (Manganese): 55mg, E6 (Zinc): 206mg.

Preservatives - Antioxidants.



Available in
2.5kg and 4.5kg bag



Approximate size

KEY VALUES

per 100g
as fed

| | |
|------------------------|---------|
| Protein | 25.5g |
| Fat content | 20g |
| Carbohydrate | 34.5g |
| NFE | 39.1g |
| Dietary fibre | 8.2g |
| Crude fibre | 3.6g |
| Omega 6 | 4.74g |
| Omega 3 | 0.83g |
| EPA+DHA | 0.32g |
| Linoleic acid | 4.56g |
| Calcium | 0.73g |
| Phosphorus | 0.7g |
| Sodium | 0.6g |
| Copper | 1.5mg |
| Zinc | 25.9mg |
| Vitamin A | 2500IU |
| Biotin | 0.33mg |
| Pantothenic acid | 14.9mg |
| Niacin | 50mg |
| Arginine | 1.84g |
| Metabolisable energy** | 410kcal |

*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation

**Calculated according to NRC2006

***NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.

CLINICAL TRIALS OR STUDIES CARRIED OUT WITH ROYAL CANIN HYPOALLERGENIC

- Mandigers PJ & al. Efficacy of a commercial hydrolysate diet in eight cats suffering from inflammatory bowel disease or adverse reaction to food. Tijdschr Diergeneeskd 2010; 135, 668-72.

ADVERSE REACTION TO FOOD



SKIN

RECOMMENDED FOR CATS WITH

- Adverse Food Reactions (AFR):
 - Food allergy
 - Food intolerance
 - With dermatological and/or gastro-intestinal signs
 - Food elimination trial
- Inflammatory Bowel Disease (IBD)
- Chronic diarrhoea
- Small intestinal bacterial overgrowth (SIBO)
- Exocrine Pancreatic Insufficiency (EPI)

NOT RECOMMENDED FOR

- Pregnancy, lactation, growth

RECOMMENDED DAILY INTAKE & FEEDING DURATION

| Cat's weight | ADULT CAT | | | | | |
|--------------|-----------|-------|--------|------|------------|------|
| | thin | | normal | | overweight | |
| | grams | cups | grams | cups | grams | cups |
| 2kg | 37 | 3/8 | 31 | 2/8 | | |
| 3kg | 50 | 4/8 | 41 | 3/8 | | |
| 4kg | 61 | 5/8 | 51 | 4/8 | 41 | 3/8 |
| 5kg | 71 | 6/8 | 59 | 5/8 | 47 | 4/8 |
| 6kg | 81 | 6/8 | 68 | 5/8 | 54 | 4/8 |
| 7kg | 90 | 7/8 | 75 | 6/8 | 60 | 5/8 |
| 8kg | 99 | 1 | 83 | 6/8 | 66 | 5/8 |
| 9kg | 108 | 1 | 90 | 7/8 | 72 | 6/8 |
| 10kg | 117 | 1 1/8 | 97 | 1 | 78 | 6/8 |

When a food allergy or a food intolerance is suspected, the cat's diet should be changed without any food transition. Nutritional management with Hypoallergenic may continue for the cat's lifetime.

It is recommended to divide the daily intake into 2 meals.



This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.