

SENSITIVITY CONTROL



- Duck and rice are the only sources of protein and carbohydrate in this diet. Both are rarely involved in food allergy, reducing the risk of GI or dermatological reactions.



- A combination of nicotinamide, inositol, choline, histidine and pantothenic acid helps reduce trans-epidermal water loss and strengthens the barrier effect of the skin.



- Eicosapentaenoic and docosahexaenoic acids (omega-3 long chain fatty acids) modulate skin reactions and contribute to the intestinal mucosal integrity.



- Contains a combination of highly digestible proteins (L.I.P.* proteins), prebiotics, beet pulp, rice and fish oil to ensure maximum digestive security.

COMPOSITION

Rice, dehydrated duck protein, vegetable fibres, hydrolysed poultry proteins, animal fats, minerals, fish oil, soya oil, marigold extract (source of lutein).

Additives (kg)***

Nutritional additives:

Vitamin A: 29700IU, Vitamin D3: 800IU, E1 (Iron): 40mg, E2 (Iodine): 3.6mg, E4 (Copper): 7mg, E5 (Manganese): 53mg, E6 (Zinc): 192mg, E8 (Selenium): 0.1mg.

Preservatives - Antioxidants.



Available in 1.5kg and 3.5kg bag



Approximate size

KEY VALUES

per 100g as fed

Protein	27g
Fat content	11g
Carbohydrate	37.6g
NFE	41.7g
Dietary fibre	10.1g
Crude fibre	6g
Omega 6	2.78g
Omega 3	0.73g
EPA+DHA	0.35g
Calcium	1.65g
Phosphorus	1.17g
Sodium	0.8g
Biotin	0.31mg
Pantothenic acid	14.64mg
Niacin	49mg
Metabolisable energy**	352kcal

*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation

**Calculated according to NRC2006

***NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet. For total levels in the diet, refer to the Average Analysis pages.

LITERATURE ON WHICH ROYAL CANIN SENSITIVITY CONTROL IS BASED

- Roudabush P. Ingredients and foods associated with adverse reactions in dogs and cats. Veterinary Dermatology 2013; 24(2): 293-294

ADVERSE REACTION TO FOOD



SKIN

RECOMMENDED FOR CATS WITH

- Adverse Food Reactions (AFR):
 - Food allergy
 - Food intolerance

NOT RECOMMENDED FOR

- Pregnancy, lactation, growth

RECOMMENDED DAILY INTAKE & FEEDING DURATION

Cat's weight	thin		normal		overweight	
	grams	cups	grams	cups	grams	cups
2kg	43	4/8	36	3/8		
3kg	58	5/8	48	4/8		
4kg	71	6/8	59	5/8		
5kg	83	7/8	69	6/8	55	5/8
6kg	95	1	79	7/8	63	5/8
7kg			88	7/8	70	6/8
8kg			97	1	77	6/8
9kg			105	1 1/8	84	7/8
10kg			113	1 2/8	91	1

When a food allergy or a food intolerance is confirmed, the cat's diet should be changed without any food transition. Continue nutritional management for the cat's lifetime.

It is recommended to divide the daily intake into 2 meals.



This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.